

Are coffee and tea consumption associated with urinary tract cancer risk? A systematic review and meta-analysis.

Citation for published version (APA):

Zeegers, M. P. A., Tan, E. S., Goldbohm, R. A., & van den Brandt, P. A. (2001). Are coffee and tea consumption associated with urinary tract cancer risk? A systematic review and meta-analysis. *International Journal of Epidemiology*, 30, 353-362. <https://doi.org/10.1093/ije/30.2.353>

Document status and date:

Published: 01/01/2001

DOI:

[10.1093/ije/30.2.353](https://doi.org/10.1093/ije/30.2.353)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Are coffee and tea consumption associated with urinary tract cancer risk? A systematic review and meta-analysis

Maurice PA Zeegers,^a Frans ES Tan,^b R Alexandra Goldbohm^c and Piet A van den Brandt^a

Background	Narrative reviews have concluded that there is a small association between coffee consumption and an increased risk of urinary tract cancer, possibly due to confounding by smoking. No association for tea consumption has been indicated. This systematic review attempts to summarize and quantify these associations both unadjusted and adjusted for age, smoking and sex.
Method	Thirty-four case-control and three follow-up studies were included in this systematic review. Summary odds ratios (OR) were calculated by meta-regression analyses.
Results	The unadjusted summary OR indicated a small increased risk of urinary tract cancer for current coffee consumers versus non-drinkers. The adjusted summary OR were: 1.26 (95% CI: 1.09–1.46) for studies with only men, 1.08 (95% CI: 0.79–1.46) for studies with only women and 1.18 (95% CI: 1.01–1.38) for studies with men and women combined. Neither unadjusted nor adjusted summary OR provided evidence for a positive association between tea consumption and urinary tract cancer. Even though studies differed in methodology, the results were rather consistent. We did not perform dose-response analyses for coffee and tea consumption due to sparse data.
Conclusions	In accordance with earlier reviews, we found that coffee consumption increases the risk of urinary tract cancer by approximately 20%. The consumption of tea seems not to be related to an increased risk of urinary tract cancer.
Keywords	Coffee, tea, urological neoplasms, bladder neoplasms, meta-analysis, epidemiology
Accepted	12 October 2000

Over the last four decades, a number of epidemiological studies have been conducted to investigate determinants of urinary tract cancer.^{1–5} These studies suggested that bladder cancer is influenced by environmental factors, such as cigarette smoking and exposure to industrial chemicals (e.g. aromatic amines), and by chronic infections with *Schistosoma haematobium*. The impact of coffee and tea consumption on the risk of urinary tract cancer is less clear, although many case-control studies and several follow-up studies on these associations have been published. Because coffee and tea are two of the most popular beverages consumed worldwide, the relationship between coffee and tea consumption and the incidence of bladder cancer is an important concern. Earlier narrative reviews on coffee consumption concluded that the evidence was consistent with a

small positive association.^{4,6–8} Viscoli *et al.*⁹ quantified this association in 1993 in a meta-analysis based on seven studies and concluded that the best available evidence does not suggest a clinically important association between coffee consumption and cancer of the lower urinary tract. According to four narrative reviews, there is no positive association between tea consumption and urinary tract cancer, although a summary association has never been quantified.^{4,6,10,11} The present study aims at reviewing all epidemiological studies up to February 2000 more systematically, to provide quantitative summary estimates of the risk of urinary tract cancer associated with both coffee and tea consumption and to evaluate changes in summary estimates after adjustment for cigarette smoking.

Material and Methods

Search strategy

The study design has been published previously.¹² Epidemiological studies were identified by the principal investigator through a computerized Medline, Cancerlit and Current Contents search on follow-up and case-control studies of all languages published

^a Department of Epidemiology, and ^b Department of Methodology and Statistics, Maastricht University, Maastricht, The Netherlands.

^c Department of Epidemiology, TNO Nutrition and Food Research Institute, Zeist, The Netherlands.

Correspondence: MPA Zeegers, Maastricht University, Department of Epidemiology, PO Box 616, 6200 MD Maastricht, The Netherlands. E-mail:mpa.zeegers@epid.unimaas.nl

until February 2000 using Medical Subjects Headings and free text words. The keywords used were urolo*, bladder, cyst*, vesic*, kidney, glomerul*, nephro*, pyel*, renal, ureteral, urethral, transitional cell, cancer, carcino*, tumo*, neoplasm*, onco*, risk, etiology, epidemiology, and caus*. Citation tracking was performed until no new studies were found. For inclusion in this analysis, the articles had to provide sufficient information to estimate a summary odds ratio (OR) and the associated standard error of incident primary urinary tract cancer for coffee or tea consumers compared to non-drinkers, i.e. exposure frequency distribution, exposure-specific OR or exposure-specific incidence rate ratios. Coffee and tea consumption were defined as ever consumption or current consumption at baseline (for follow-up studies) or in the reference period (for case-control studies).

Data collection

A criteria list for the assessment of quality items (study characteristics) in observational cancer research was developed (Appendix 1). The criteria list was used to provide co-variables for meta-regression to explore reasons for observed heterogeneity in results and is available upon request. The criteria list was developed through consensus meetings with experts in the fields of cancer and meta-analysis. The list comprised items felt to be important for the quality of observational research, including study design (follow-up or case control study), measuring instrument (interview or self-administered questionnaire) and anatomical site of the neoplasm (total urinary tract, bladder, renal pelvis or ureter). For case-control studies, additional information was gathered on the sources of the cases and controls. Case-control studies that used controls obtained from the general population or from hospitals were defined as population-based or hospital-based case control studies, respectively. If case-control studies used both population- and hospital-based controls, population-based controls were used for further analysis. Data were extracted to permit the calculation of both unadjusted and adjusted OR to estimate the association between coffee and tea consumption and the risk of urinary tract cancer. Two-way contingency tables for each study were constructed, based on exposure frequency distributions, in order to calculate the unadjusted OR. Adjusted OR were extracted directly from the original reports. Because we considered age, smoking and sex to be the most important confounding variables, the authors of the original articles had to have adjusted for at least these three variables for inclusion in the calculation of adjusted summary estimates. If studies reported sex-stratified age- and smoking-adjusted OR, we combined these estimates by calculating age-, smoking- and sex-adjusted OR.¹³ In such calculations, the inverse of the variance was used to weight the reported sex-stratified estimates. For studies that reported separate adjusted OR for several consumption strata, we estimated a total OR for current or ever use weighted by the exposure specific prevalence of the non-cases.¹⁴

Statistical analysis

To detect publication or related biases, we explored asymmetry in funnel plots, i.e. plots of effect estimates against their estimated precision.¹⁵ This is examined visually and the degree of asymmetry is measured using Egger's unweighted regression asymmetry test.^{16,17} If a study appeared in more than one publication, data from the last publication were used for statistical analysis. We estimated the summary OR and corresponding

95% CI with random effects meta regression analysis using the Stata statistical software.^{18,19} The between-study variance was estimated iteratively, using the empirical Bayes method.^{19,20} Summary OR were calculated for total coffee consumption (current drinker versus non-drinker and ever drinker versus never-drinker), decaffeinated coffee (drinker versus non-drinker) and total tea consumption (current drinker versus non-drinker and ever drinker versus never-drinker). We did not perform subgroup analyses by beverage type other than decaffeinated coffee or by the number of cups consumed per day due to sparse data. Data were analysed separately or combined for men and women, depending on data availability in the original studies. Furthermore, we evaluated the influence of the type of adjustment for cigarette smoking in the component studies on the summary association between current coffee and tea consumption and bladder cancer risk. This analysis might give some insight into the existence of residual confounding from inadequate adjustment of cigarette smoking in the component studies. To explore reasons for observed heterogeneity, sensitivity analyses were performed on the study characteristics measuring instrument, anatomical site of the neoplasm and source of cases and controls, and their influence on the association between coffee or tea consumption and urinary tract cancer was tested.

Results

Study characteristics

We identified 69 articles reporting follow-up or case-control studies on coffee and tea consumption and urinary tract cancer published between 1968 and 1999.²¹⁻⁸⁹ Generally, the association between coffee or tea consumption and urinary tract cancer was not the main research hypothesis. Twenty articles were excluded because they did not provide sufficient information to estimate a summary OR and corresponding standard error^{21-29,33-39} or did not use incident cases.^{30-32,87} The remaining 49 articles described three follow-up studies^{40,41,84,88} and 34 case-control studies with incident cases.^{42-83,85,86,89} One case-control study that provided separate associations for parts of the study performed in the US, the UK and Japan was considered as three separate studies.⁴⁵ The case-control studies were population-based ($n = 12$),^{42-56,83,89} hospital-based ($n = 18$)^{57-78,85,86} or neighbourhood-based ($n = 3$).⁷⁹⁻⁸¹ One case-control study used both population- and hospital-based controls.⁸² The hospital controls in most case-control studies did not have any diet- or smoking-related disease. The case-control studies also varied with regard to their criteria of case selection. Seven case-control studies identified cancer cases in defined populations,^{44,46,47,49,51,52,55,56,79,80,83,89} 22 case-control studies selected cases from hospitals^{42,43,45,53,58-63,65-78,81,82,85,86} and five case-control studies used both sources.^{48,50,54,57,64} Information on coffee and tea consumption was obtained by interview,^{40,42-58,61,62,64-83,86,89} self-administered questionnaire^{41,59,60,85,88} or both techniques.^{63,84} Some studies included all neoplasms of the urinary tract as cases, of which more than 90% were found to involve bladder cancer,^{42,45,48,53,60,73,84,85} while others selected only bladder carcinomas,^{40,41,44,46,47,49-52,54-59,62-72,74-79,81-83,86,88} carcinomas of the renal pelvis^{61,89} or carcinomas of renal pelvis and ureter combined.⁸⁰ Most studies used histologically confirmed cases with urothelial cell cancer (Table 1).

Table 1 Study characteristics of published epidemiological studies concerning coffee and tea consumption and cancer of the urinary tract, ordered by year of publication

Ref.	First author	Publication year	Country	Anatomical site of urinary tract	Study design		Case-control study		Coffee/Tea assessment
					Follow-up study		Case source	Control source	
57	Dunham	1968	US	bladder	–		both	hospital	interview
42	Cole ^a	1971	US	urinary tract	–		hospital	population	interview
58	Bross	1973	US	bladder	–		hospital	hospital	interview
59	Morgan	1974	Canada	bladder	–		hospital	hospital	questionnaire
43	Schmauz ^a	1974	US	urinary tract	–		hospital	population	interview
60	Simon	1975	US	urinary tract	–		hospital	hospital	questionnaire
61	Armstrong	1976	UK	renal pelvis	–		hospital	hospital	interview
44	Miller ^b	1977	Canada	bladder	–		population	population	interview
62	Wynder ^c	1977	US	bladder	–		hospital	hospital	interview
63	Mettlin	1979	US	bladder	–		hospital	hospital	both
79	Howe ^b	1980	Canada	bladder	–		population	neighbour	interview
64	Cartwright	1981	UK	bladder	–		both	hospital	interview
45	Morrison	1982	US	urinary tract	–		hospital	population	interview
45	Morrison	1982	UK	urinary tract	–		hospital	population	interview
45	Morrison ^d	1982	Japan	urinary tract	–		hospital	population	interview
65	Najem	1982	US	bladder	–		hospital	hospital	interview
46	Hartge ^e	1983	US	bladder	–		population	population	interview
47	Marrett ^e	1983	US	bladder	–		population	population	interview
89	McLaughlin	1983	US	renal pelvis	–		population	population	interview
48	Ohno ^d	1985	Japan	urinary tract	–		both	population	interview
66	Rebekalos	1985	Greece	bladder	–		hospital	hospital	interview
85	Bravo	1986	Spain	urinary tract	–		hospital	hospital	questionnaire
40	Heilbrun ^f	1986	US	bladder	yes		–	–	interview
49	Jensen	1986	Denmark	bladder	–		population	population	interview
67	Kabat ^c	1986	US	bladder	–		hospital	hospital	interview
83	Piper	1986	US	bladder	–		population	population	interview
68	Iscovich	1987	US	bladder	–		hospital	hospital	interview
69	Ciccone	1988	Italy	bladder	–		hospital	hospital	interview
50	Risch	1988	Canada	bladder	–		both	population	interview
52	Slattery ^e	1988	US	bladder	–		population	population	interview
51	Slattery ^e	1988	US	bladder	–		population	population	interview
70	LaVecchia ^g	1989	Italy	bladder	–		hospital	hospital	interview
80	Ross	1989	US	renal pelvis ^h	–		population	neighbour	interview
71	Clavel	1991	France	bladder	–		hospital	hospital	interview
41	Mills	1991	US	bladder	yes		–	–	questionnaire
53	Nomura	1991	US	urinary tract	–		hospital	population	interview
72	d'Avanzo ^g	1992	Italy	bladder	–		hospital	hospital	interview
73	Kunze	1992	Germany	urinary tract	–		hospital	hospital	interview
74	LaVecchia ^g	1992	Italy	bladder	–		hospital	hospital	interview
84	Chyou ^f	1993	US	urinary tract	yes		–	–	both
82	Pujolar	1993	Spain	bladder	–		hospital	both	interview
81	Vena	1993	US	bladder	–		hospital	neighbour	interview
75	Barbone ^g	1994	Italy	bladder	–		hospital	hospital	interview
76	Hours	1994	France	bladder	–		hospital	hospital	interview
54	Momas	1994	France	bladder	–		both	population	interview
55	Sturgeon ^e	1994	US	bladder	–		population	population	interview
77	d'Avanzo ^g	1995	Italy	bladder	–		hospital	hospital	interview
56	Bruemmer	1997	US	bladder	–		population	population	interview
78	Donato ^h	1997	Italy	bladder	–		hospital	hospital	interview
86	Donato ^h	1998	Italy	bladder	–		hospital	hospital	interview
88	Michaud	1999	US	bladder	yes		–	–	questionnaire

a,b,c,d,e,f,g same study has appeared in more than one publication, ^h and ureter

Risk estimation

Funnel plots showed no asymmetry, neither visually nor in terms of statistical significance (P -values ≥ 0.51 and ≥ 0.12 for current coffee and tea consumption, respectively) (Figure 1). Table 2 summarizes the unadjusted and adjusted results of observational studies reporting the association between coffee and tea consumption. Almost all included articles provided some information to estimate an association between coffee consumption and urinary tract cancer.^{41-73,75-86,88,89} The study specific effect estimates are available upon request. The unadjusted summary OR for current coffee drinkers compared to non-drinkers were 1.30 (95% CI: 1.15–1.47) for 25 studies with only men, 1.18 (95% CI: 0.96–1.44) for 23 studies with only women and 1.26 (95% CI: 1.11–1.43) for 25 studies with men and women combined. The adjusted summary OR for current coffee consumption were somewhat lower after adjustment for age, smoking and sex. The adjusted summary OR for current coffee consumption were 1.26 (95% CI: 1.09–1.46), 1.08 (95% CI: 0.79–1.46) and 1.18 (95% CI: 1.01–1.38) for 16 studies with men, 12 studies with women and 14 studies with men and women combined. The adjusted summary associations between ever versus never coffee consumption and urinary tract cancer also showed small increased risks, although the unadjusted summary estimates were higher. The consumption of decaffeinated coffee also appeared to be associated with bladder cancer risk, although it was not statistically significant (Table 2).

Twenty articles provided sufficient information to estimate a summary association of tea consumption.^{40,46,48-53,56,59,60,68,71-74,79,84,88,89} Neither the unadjusted nor the age-, smoking-, and sex-adjusted summary associations provided evidence for a positive association between current tea consumption and the risk of urinary tract cancer (Table 2). The sex-specific adjusted summary OR for current tea consumers compared to non-drinkers were 1.08 (95% CI: 0.94–1.24) for seven studies with men and 0.99 (95% CI: 0.81–1.20) for six studies with women. Seven studies allowed calculation of adjusted summary OR for men and women combined: 1.01 (95% CI: 0.92–1.10). Very

few studies reported an effect estimate for ever versus never tea consumption. Both unadjusted and adjusted summary associations for ever tea consumption did not indicate a positive association between tea consumption and bladder cancer risk. All summary associations were around unity (Table 2).

Confounding by cigarette smoking

Seventeen studies presented adjusted OR for current coffee and tea consumption for men and women combined. Of these, seven studies adjusted for cigarette smoking by incorporating one smoking variable (mostly smoking status) in a regression model.^{48,56,62,65,69,77,82} Seven other studies used a combination of cigarette smoking features (mostly amount and duration) to adjust for cigarette smoking.^{49,50,52,53,55,73,78} The type of adjustment for three studies remains unclear.^{41,66,74} The summary OR for current coffee consumption compared to non-drinkers was larger in studies in which several cigarette smoking features were used to adjust for cigarette smoking compared to studies in which only one smoking variable was used (Table 3). For tea consumption, the type of adjustment for cigarette smoking did not influence the summary associations substantially (Table 3).

Sensitivity analysis

We further examined the crude association of current coffee and tea consumption and urinary tract cancer by measuring instrument, sources of cases and controls and anatomical site of the tumour to explore their influence on the outcome estimates in case-control studies that provided information for men and women combined (Figure 2). Unfortunately, none of the follow-up studies provided sufficient information for men and women combined to contribute to this sensitivity analysis. All tests for multiplicative interaction were statistically non-significant. Most subset-specific summary OR did not differ substantially. Neither selection on anatomical site of the tumour nor selection on the source of cases and controls altered the summary OR. It appeared, however, that the summary associations for studies that used interviewing techniques were higher than for studies that used self-administered questionnaires (Figure 2).




Figure 1 Funnel plots for current coffee and tea drinkers versus non-drinkers, unadjusted. Interrupted and uninterrupted reference lines indicate total summary odds and null effect.

Table 2 Summary odds ratios (OR) for coffee and tea consumption both unadjusted and adjusted for age, smoking and sex calculated from epidemiological studies

Coffee/Tea consumption	Male			Female			Male and Female		
	No. ^a	OR (95% CI)	Q ^b (P)	No. ^a	OR (95% CI)	Q ^b (P)	No. ^a	OR (95% CI)	Q ^b (P)
Unadjusted summary odds ratios									
Coffee consumption									
Current versus non drinker	25	1.30 (1.15–1.47)	45.94 (<0.01)	23	1.18 (0.96–1.44)	35.00 (0.04)	25	1.26 (1.11–1.43)	56.36 (<0.01)
Ever versus never drinker	7	1.92 (1.37–2.70)	13.57 (0.04)	6	1.66 (0.81–3.38)	15.47 (<0.01)	7	1.85 (1.21–2.82)	22.43 (<0.01)
Decaffeinated coffee consumption									
Drinker versus non-drinker	4	1.08 (0.88–1.32)	3.79 (0.29)	2	0.56 (0.35–0.91)	0.01 (0.91)	4	1.01 (0.74–1.37)	6.52 (0.09)
Tea consumption									
Current versus non-drinker	8	1.04 (0.89–1.21)	10.11 (0.18)	8	0.97 (0.78–1.20)	9.179 (0.24)	0.94	14.59 (0.82–1.08)	(0.07)
Ever versus never drinker	3	0.94 (0.66–1.35)	3.69 (0.16)	2	1.36 (0.18–10.11)	14.90 (<0.01)	3	0.77 (0.47–1.25)	7.16 (0.03)
Adjusted summary odds ratios									
Coffee consumption									
Current versus non-drinker	16	1.26 (1.09–1.46)	40.15 (<0.01)	12	1.08 (0.79–1.46)	26.68 (<0.01)	14	1.18 (1.01–1.38)	30.03 (<0.01)
Ever versus never drinker	6	1.19 (1.00–1.41)	6.03 (0.30)	5	1.16 (0.86–1.56)	4.25 (0.37)	5	1.18 (1.03–1.36)	1.06 (0.90)
Decaffeinated coffee consumption									
Drinker versus non-drinker	5	1.04 (0.86–1.25)	4.47 (0.35)	3	1.18 (0.63–2.22)	5.76 (0.06)	4	1.18 (0.99–1.40)	2.40 (0.49)
Tea consumption									
Current versus non-drinker	7	1.08 (0.94–1.24)	9.07 (0.17)	6	0.99 (0.81–1.20)	6.96 (0.22)	7	1.01 (0.92–1.10)	5.47 (0.49)
Ever versus never drinker	4	1.02 (0.84–1.25)	2.97 (0.40)	3	1.00 (0.27–3.73)	13.81 (<0.01)	3	1.09 (0.54–2.17)	9.60 (0.15)

^a Number of analysed studies.^b χ^2 -test for heterogeneity.**Table 3** Summary odds ratios (OR) for current coffee and tea consumption compared to non-drinkers according to the type of adjustment for cigarette smoking made in the component studies, for men and women combined

Coffee/tea consumption	Adjusted for one cigarette smoking variable ^a			Adjusted for several cigarette smoking features ^b		
	N ^c	OR (95% CI)	Q ^d (P)	N ^c	OR (95% CI)	Q ^d (P)
Coffee consumption						
Current versus non-drinker	7	1.04 (0.89–1.22)	8.90 (0.18)	5	1.36 (1.08–2.01)	11.74 (0.02)
Tea consumption						
Current versus non-drinker	1	1.02 (0.77–1.33)	N/A	5	1.09 (0.96–1.25)	3.25 (0.52)

^a Adjusted for cigarette smoking status or amount only.^b Adjusted for any combination of cigarette smoking status, amount or duration.^c Number of analysed studies.^d χ^2 -test for heterogeneity.




Figure 2 Forest plots of crude summary odds ratios and 95% confidence intervals of urinary tract cancer for current coffee and tea consumers compared to non-drinkers, by measuring instrument, source of cases and controls and tumour site for case-control studies with men and women combined

Interrupted and uninterrupted reference lines indicates total summary odds and null effect.

Discussion

We were able to retrieve 49 articles reporting follow-up or case-control studies on coffee and tea consumption and the risk of human urinary tract cancer in the present meta-analysis. These observational studies can be considered as the best available epidemiological evidence, because no human experimental studies have been conducted. The available evidence suggests a small increased risk for coffee consumers remaining after adjustment for age, smoking and sex. The consumption of tea seems not to be related to an increased risk of urinary tract cancer.

We did not attempt to uncover unpublished observations and excluded studies that did not meet the predetermined criteria. The excluded studies showed inconsistent results for coffee consumption. The estimated associations of coffee consumption ranged from moderately inverse,^{27,32,38} through no association^{24–26,28,29,31,90} to moderately positive.^{22,23,30,33–37,39,91} Although three of the excluded studies on tea consumption have presented positive associations,^{34,37} most studies could not confirm these findings.^{21,23,27,33,35,90} Publication bias might arise by excluding these studies. However, we did not identify funnel plot asymmetry in our meta-analysis, either visually or in terms of statistical significance.

The definition of 'current drinker' at baseline (follow-up studies) or in the reference period (case-control studies) might have caused heterogeneity between studies, because the follow-up period and the reference date varied between the included studies. Therefore, results have to be interpreted with caution. Because of potential additional heterogeneity in populations, designs and analyses of various studies, we assumed that the true effects being estimated would vary between the studies in addition to the usual sampling variation in the estimates (within studies), even if the test for heterogeneity could not identify heterogeneity. To account for both sources of variation we used a random effects meta regression analysis to combine the

results from the primary studies.²⁰ The random effect approach provides some allowance for heterogeneity in studies beyond sampling error.

The summary OR were similar across sources of the cases and controls in the case-control studies. It appeared that the anatomical site of the tumour did not modify the association between coffee/tea consumption and urinary tract cancer risk. Because the overwhelming majority of urinary tract tumours occurred in the urinary bladder and, since the renal pelvis and urether are covered by the same urothelium as the urinary bladder, no biologically plausible difference would be expected. Although we could not find statistically significant interaction effects, it appeared that the summary estimates of coffee and tea consumption were higher for studies that used interviewing techniques than for studies that used self-administered questionnaires. This contrast can be a consequence of response bias due to different assessment techniques or to chance alone.

Limitations in reported data meant that the summary OR could not be quantified by amount or type of tea or coffee, other than decaffeinated coffee. The included studies provided data on coffee or tea consumption within exposure categories with different cut-off points. Therefore, we had to dichotomize coffee and tea consumption in current consumption versus no consumption as well as ever consumption versus never consumption. It should be noted, that a dose-response analysis would give more information about the underlying association of coffee and tea consumption^{92,93} with urinary tract cancer. However, the result of a dose-response analysis appears to depend heavily on the choice of assigned dose of the different exposure categories.⁹² Hence, for a proper dose-response analysis, the number of exposure levels should be at least equal to four. More than 25% of the component studies reported three exposure levels at the most for coffee^{55,57,63–66,73,76,77,84,85,87} and tea consumption.^{48,50,52,53,72,74,79,73} Moreover, for more than 25% of the component studies, no dose-response analysis for the adjusted

OR was possible, because no exposure specific confidence intervals were reported. In fact, in only 14 out of 46 studies on coffee consumption and in two out of 14 on tea consumption, was a dose response analysis tolerable. With the above-mentioned considerations in mind, we have decided not to conduct dose-response analyses at all.

Although 49 articles contributed to this meta-analysis, only 30 provided sufficient information to permit calculation of OR, adjusted for at least age, smoking and sex. It was not possible to adjust for a specific set of confounders. The decision of authors to report adjusted effect estimates or not, may have been influenced by the results of the component studies. Similarly, the type of confounders that authors chose to adjust for varied and may to some extent have been driven by the data. Therefore, we also calculated unadjusted OR. Furthermore, focussing entirely on adjusted OR would have resulted in the exclusion of 39% of the studies that also had (crude) information on the association between coffee or tea consumption and cancer of the urinary tract. The presentation of both unadjusted and adjusted OR might give some insight into the potential confounding effect in particular of cigarette smoking. It remains possible that adjustment for smoking in the component studies, and therefore in our summary estimates, may have been inadequate if smoking was not carefully adjusted for. Yet, it appeared that adjustment for several cigarette smoking features in the component studies, compared to adjustment for one smoking variable alone, was not associated with a decreased risk of bladder cancer, as would be expected in the case of residual confounding. However, these results should be carefully interpreted since the comparisons

were based on different subsets of studies. To avoid residual confounding, the impact of coffee drinking on bladder cancer risk can be evaluated in lifelong non-smokers.⁷ Of all identified articles, four reported no increased risk associated with coffee drinking among non-smokers^{23,58,67,79}, whereas 11 studies suggested a moderately increased risk.^{41,46,50,51,58,60,66,69,71,72,82}

In addition to residual confounding by cigarette smoking, another explanation for a small positive association between coffee consumption and cancer of the urinary tract might be that non-coffee-drinkers are a rather selected population. Non-coffee drinkers may differ also in occupation, the consumption of alcohol, tea, vegetables or fruit, or other lifestyle habits from the general population of coffee drinkers.

The consumption of tea seems not to be related to an increased risk of urinary tract cancer, although the results are based on only a few studies. Evidence from animal experimental studies shows that it is plausible that certain compounds present in tea (e.g. flavonoids) may even protect against the development of cancer.⁹⁴ If an inverse association between tea consumption and urinary tract cancer does exist, it is likely to be small. Further research from large epidemiological studies is needed in this area.

In accordance with earlier reviews and based on predominantly case-control data, we found that coffee consumption increases the risk of urinary tract cancer by approximately 20%. The consumption of tea seems not to be related to an increased risk of urinary tract cancer. The risk of urinary tract cancer related to the amount or type of coffee and tea needs further investigation.

KEY MESSAGES

- Based on a meta-analysis of 37 epidemiological studies, we estimated that coffee consumption increases urinary tract cancer risk by approximately 20%.
- The consumption of tea seems not to be related to an increased risk of urinary tract cancer.

References

- Silverman DT, Hartge P, Morrison AS *et al.* Epidemiology of bladder cancer. *Hematol Oncol Clin North Am* 1992;**6**:1–30.
- Johansson SL, Cohen SM. Epidemiology and etiology of bladder cancer. *Semin Surg Oncol* 1997;**13**:291–98.
- Ross RK, Jones PA, Yu MC. Bladder cancer epidemiology and pathogenesis. *Semin Oncol* 1996;**23**:536–45.
- World Cancer Research Fund, American Institute for Cancer Research. Bladder. In: *Food, Nutrition and the Prevention of Cancer: A Global Perspective*. Menasha: Banta Book Group, 1997, pp.338–61.
- van der Meijden APM. Bladder cancer. *Br Med J* 1998;**317**:1366–69.
- WHO, International Agency for Research on Cancer. *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans: Coffee Tea, Mate, Methylxanthines and Methylglyoxal*, Vol. 51. Lyon: World Health Organization, 1991.
- Silverman DT, Morrison AS, Devesa SS. Bladder cancer. In: Schottenfeld D, Fraumeni JF (eds). *Cancer Epidemiology and Prevention* New York Oxford: Oxford University Press, 1996, pp.1156–79.
- Nehlig A, Debry G. Coffee and cancer: a review of human and animal data. *World Rev Nutr Diet* 1996;**79**:185–221.
- Viscoli CM, Lachs MS, Horwitz RJ. Bladder cancer and coffee drinking: a summary of case-control research. *Lancet* 1993;**341**:1432–37.
- Blot WJ, Chow WH, McLaughlin JK. Tea and cancer: a review of the epidemiological evidence. *Eur J Cancer Prev* 1996;**5**:425–38.
- Katiyar SK, Mukhtar H. Tea consumption and cancer. *World Rev Nutr Diet* 1996;**79**:154–84.
- Zeegers MPA, Tan FES, Verhagen AP *et al.* Elevated risk of cancer of the urinary tract for alcohol drinkers: a meta-analysis. *Cancer Causes Control* 1999;**10**:445–51.
- Woolf B. On estimation of the relation between blood group and disease. *Ann Hum Genet* 1954;**19**:251–53.
- Tan FES, Zeegers MPA. An alternative method of combining exposure specific odds ratios in epidemiologic studies. In: *Research Report*. Maastricht: Maastricht University. Department Methodology and Statistics, 1999.
- Light RJ, Pillemer DB. Quantitative procedures. In: *Summing Up: The Science of Reviewing Research*. Harvard: Harvard University Press, 1984.
- Egger M, Davey Smith G, Schneider M *et al.* Bias in meta-analysis detected by a simple, graphical test. *Br Med J* 1997;**315**:629–34.
- Steichen TJ, Egger M, Sterne J. Tests for publication bias in meta-analysis. *Stata Tech Bull* 1998;**STB-44**:sbe 19.1.

- 18 StataCorp. *Stata Statistical Software: Release 6.0*. College Station, TX: Stata Corporation, 1999.
- 19 Sharp SJ. Meta-analysis regression. *Stata Tech Bull* 1998;**42**: sbe 23.
- 20 Berkey CS, Hoaglin DC, Mosteller F *et al*. A random-effects regression model for meta-analysis. *Stat Med* 1995;**14**:395-411.
- 21 Zheng W, Doyle TJ, Kushi LH *et al*. Tea consumption and cancer incidence in a prospective cohort study of postmenopausal women. *Am J Epidemiol* 1996;**144**:175-82.
- 22 Bravo MP, Del Rey Calero J, Conde M. Risk factors of bladder cancer in Spain. *Neoplasma* 1987;**34**:633-37.
- 23 Akdas A, Kirkali Z, Bilir N. Epidemiological case-control study on the etiology of bladder cancer in Turkey. *Eur Urol* 1990;**17**:23-26.
- 24 Mommsen S, Aagaard J, Sell A. An epidemiological study of bladder cancer in a predominantly rural district. *Scand J Urol Nephrol* 1983;**17**:307-12.
- 25 Mommsen S, Aagaard J, Sell A. An epidemiological case-control study of bladder cancer in males from a predominantly rural district. *Eur J Cancer Clin Oncol* 1982;**18**:1205-10.
- 26 Mommsen S, Aagaard J. A case-control study of bladder-cancer: a multivariate, stratified analysis of a low-risk population. *Dan Med Bull* 1983;**30**:427-32.
- 27 Whittemore AS, Paffenbarger RS Jr, Anderson K *et al*. Early precursors of urogenital cancers in former college men. *J Urol* 1984;**132**: 1256-61.
- 28 Wynder EL, Dieck GS, Hall NE *et al*. A case-control study of diesel exhaust exposure and bladder cancer. *Environ Res* 1985;**37**:475-89.
- 29 Whittemore AS, Paffenbarger RS Jr, Anderson K *et al*. Early precursors of site-specific cancers in college men and women. *J Natl Cancer Inst* 1985;**74**:43-51.
- 30 Snowdon DA, Phillips RL. Coffee consumption and risk of fatal cancers. *Am J Public Health* 1984;**74**:820-23.
- 31 Kinlen LJ, Willows AN, Goldblatt P *et al*. Tea consumption and cancer. *Br J Cancer* 1988;**58**:397-401.
- 32 Gonzalez CA, Lopez Abente G, Errezola M *et al*. Occupation, tobacco use, coffee, and bladder cancer in the county of Mataro (Spain). *Cancer* 1985;**55**:2031-34.
- 33 Miller AB. The etiology of bladder cancer from the epidemiological viewpoint. *Cancer Res* 1977;**37**:2939-42.
- 34 Miller CT, Neutel CI, Nair RC *et al*. Relative importance of risk factors in bladder carcinogenesis. *J Chron Dis* 1978;**31**:51-56.
- 35 Sullivan JW. Epidemiologic survey of bladder cancer in greater New Orleans. *J Urol* 1982;**128**:281-83.
- 36 Mommsen S, Aagaard J, Sell A. A case-control study of female bladder cancer. *Eur J Cancer Clin Oncol* 1983;**19**:725-29.
- 37 Claude J, Kunze E, Frentzel Beyne R *et al*. Life-style and occupational risk factors in cancer of the lower urinary tract. *Am J Epidemiol* 1986;**124**:578-89.
- 38 Jacobsen BK, Bjelke E, Kvale G *et al*. Coffee drinking, mortality, and cancer incidence: results from a Norwegian prospective study. *J Natl Cancer Inst* 1986;**76**:823-31.
- 39 Fraumeni JF Jr, Scotto J, Dunham LJ. Coffee-drinking and bladder cancer. *Lancet* 1971;**ii**:1204.
- 40 Heilbrun LK, Nomura A, Stemmermann GN. Black tea consumption and cancer risk: a prospective study. *Br J Cancer* 1986;**54**:677-83.
- 41 Mills PK, Beeson WL, Phillips RL *et al*. Bladder cancer in a low risk population: results from the Adventist Health Study. *Am J Epidemiol* 1991;**133**:230-39.
- 42 Cole P. Coffee-drinking and cancer of the lower urinary tract. *Lancet* 1971;**26**:1335-37.
- 43 Schmauz R, Cole P. Epidemiology of cancer of the renal pelvis and urether. *J Natl Cancer Inst* 1974;**52**:1431-34.
- 44 Miller AB, Howe GR. Artificial sweeteners and bladder cancer. *Lancet* 1977;**10**:1221-22.
- 45 Morrison AS, Buring JE, Verhoek WG *et al*. Coffee drinking and cancer of the lower urinary tract. *J Natl Cancer Inst* 1982;**68**:91-94.
- 46 Hartge P, Hoover R, West DW *et al*. Coffee drinking and risk of bladder cancer. *J Natl Cancer Inst* 1983;**70**:1021-26.
- 47 Marrett LD, Walter SD, Meigs JW. Coffee drinking and bladder cancer in Connecticut. *Am J Epidemiol* 1983;**117**:113-27.
- 48 Ohno Y, Aoki K, Obata K *et al*. Case-control study of urinary bladder cancer in metropolitan Nagoya. *Natl Cancer Inst Monogr* 1985;**69**: 229-34.
- 49 Jensen OM, Wahrendorf J, Knudsen JB *et al*. The Copenhagen case-control study of bladder cancer. II. Effect of coffee and other beverages. *Int J Cancer* 1986;**37**:651-57.
- 50 Risch HA, Burch JD, Miller AB *et al*. Dietary factors and the incidence of cancer of the urinary bladder. *Am J Epidemiol* 1988;**127**:1179-91.
- 51 Slattery ML, Schumacher MC, West DW *et al*. Smoking and bladder cancer. The modifying effect of cigarettes on other factors. *Cancer* 1988;**61**:402-08.
- 52 Slattery ML, West DW, Robison LM. Fluid intake and bladder cancer in Utah. *Int J Cancer* 1988;**42**:17-22.
- 53 Nomura AMY, Kolonel LN, Hankin JH *et al*. Dietary factors in cancer of the urinary tract. *Int J Cancer* 1991;**48**:199-205.
- 54 Momas I, Daurès JP, Festy B *et al*. Relative importance of risk factors in bladder carcinogenesis: some new results about Mediterranean habits. *Cancer Causes Control* 1994;**5**:326-32.
- 55 Sturgeon SR, Hartge P, Silverman DT *et al*. Associations between bladder cancer risk factors and tumor stage and grade at diagnosis. *Epidemiology* 1994;**5**:218-25.
- 56 Bruemmer B, White E, Vaughan TL *et al*. Fluid intake and the incidence of bladder cancer among middle-aged men and women in a three-county area of western Washington. *Nutr Cancer* 1997;**29**:163-68.
- 57 Dunham LJ, Rabson AS, Steward HS *et al*. Rates, interview, and pathology study of cancer of the urinary bladder in New Orleans, Louisiana. *J Natl Cancer Inst* 1968;**41**:683-709.
- 58 Bross IDJ, Tindings J. Another look at coffee drinking and cancer of the urinary bladder. *Prev Med* 1973;**2**:445-51.
- 59 Morgan RW, Meera GJ. Bladder cancer: smoking, beverages and artificial sweeteners. *Can Med Assoc J* 1974;**111**:1067-70.
- 60 Simon D, Yen S, Cole P. Coffee drinking and cancer of the lower urinary tract. *J Natl Cancer Inst* 1975;**54**:587-91.
- 61 Armstrong B, Garrod A, Doll R. A retrospective study of renal cancer with special reference to coffee and animal protein consumption. *Br J Cancer* 1976;**33**:127-36.
- 62 Wynder EL, Goldsmith R. The epidemiology of bladder cancer: a second look. *Cancer* 1977;**40**:1246-68.
- 63 Mettlin C, Graham S. Dietary risk factors in human bladder cancer. *Am J Epidemiol* 1979;**110**:255-63.
- 64 Cartwright RA, Adib R, Ghashan R *et al*. The epidemiology of bladder cancer in West Yorkshire: a preliminary report on non-occupational aetiologies. *Carcinogenesis* 1981;**2**:343-47.
- 65 Najem GR, Louri DB, Seebode JJ *et al*. Life time occupation, smoking, caffeine, saccharine, hair dyes and bladder carcinogenesis. *Int J Epidemiol* 1982;**11**:212-17.
- 66 Rebekalos A, Trichopoulos D, Tzonou A *et al*. Tobacco smoking, coffee drinking, and occupation as risk factors for bladder cancer in Greece. *J Natl Cancer Inst* 1985;**75**:455-61.
- 67 Kabat GC, Dieck GS, Wynder EL. Bladder cancer in nonsmokers. *Cancer* 1986;**57**:362-67.
- 68 Iscovich J, Castelletto R, Esteve J *et al*. Tobacco smoking, occupational exposure and bladder cancer in Argentina. *Int J Cancer* 1987;**40**: 734-40.

- ⁶⁹ Ciccone G, Vineis P. Coffee drinking and bladder cancer. *Cancer Lett* 1988;**41**:45-52.
- ⁷⁰ La Vecchia C, Negri E, Decarli A *et al*. Dietary factors in the risk of bladder cancer. *Nutr Cancer* 1989;**12**:93-101.
- ⁷¹ Clavel J, Cordier S. Coffee consumption and bladder cancer risk. *Int J Cancer* 1991;**47**:207-12.
- ⁷² D'Avanzo B, La Vecchia C, Franceschi S *et al*. Coffee consumption and bladder cancer risk. *Eur J Cancer* 1992;**28a**:1480-84.
- ⁷³ Kunze E, Chang Claude J, Frentzel Beyme R. Life style and occupational risk factors for bladder cancer in Germany. A case-control study. *Cancer* 1992;**69**:1776-90.
- ⁷⁴ La Vecchia C, Negri E, Franceschi S *et al*. Tea consumption and cancer risk. *Nutr Cancer* 1992;**17**:27-31.
- ⁷⁵ Barbone F, Franceschi S, Talamini R *et al*. Occupation and bladder cancer in Pordenone (north-east Italy): a case-control study. *Int J Epidemiol* 1994;**23**:58-65.
- ⁷⁶ Hours M, Danache B, Fevotte J *et al*. Bladder cancer and occupational exposures. *Scan J Work Environ Health* 1994;**20**:322-30.
- ⁷⁷ D'Avanzo B, La Vecchia C, Negri E *et al*. Attributable risks for bladder cancer in northern Italy. *Ann Epidemiol* 1995;**5**:427-31.
- ⁷⁸ Donato F, Boffetta P, Fazioli R *et al*. Bladder cancer, tobacco smoking, coffee and alcohol drinking in Brescia northern Italy. *Eur J Epidemiol* 1997;**13**:795-800.
- ⁷⁹ Howe GR, Burch JD, Miller AB *et al*. Tobacco use, occupation, coffee, various nutrients, and bladder cancer. *J Natl Cancer Inst* 1980;**64**:701-13.
- ⁸⁰ Ross RK, Paganini Hill A, Landolph J *et al*. Analgesics, cigarette smoking, and other risk factors for cancer of the renal pelvis and ureter. *Cancer Res* 1989;**49**:1045-48.
- ⁸¹ Vena JE, Freudenheim J, Graham S *et al*. Coffee, cigarette smoking, and bladder cancer in western New York. *Ann Epidemiol* 1993;**3**:586-91.
- ⁸² Pujolar EA, Gonzalez CA, Lopez Abente G *et al*. Bladder cancer and coffee consumption in smokers and non-smokers in Spain. *Int J Epidemiol* 1993;**22**:38-44.
- ⁸³ Piper JM, Matanoski GM, Tonascia J. Bladder cancer in young women. *Am J Epidemiol* 1986;**123**:1033-42.
- ⁸⁴ Chyou PH, Nomura AM, Stemmermann GN. A prospective study of diet, smoking, and lower urinary tract cancer. *Ann Epidemiol* 1993;**3**:211-16.
- ⁸⁵ Bravo P, del Rey J, Sanchez J *et al*. Cafe y analgesicos como factores de riesgo del cancer de vejiga. *Arch Esp Urol* 1986;**39**:337-41.
- ⁸⁶ Donato F, Boffetta P, Fazioli R *et al*. Reliability of data on smoking habit and coffee drinking collected by personal interview in a hospital-based case-control study. *Eur J Epidemiol* 1998;**14**:259-67.
- ⁸⁷ Stensvold I, Jacobsen BK. Coffee and cancer: a prospective study of 43 000 Norwegian men and women. *Cancer Causes Control* 1994;**5**:401-08.
- ⁸⁸ Michaud DS, Spiegelman D, Clinton SK *et al*. Fluid intake and the risk of bladder cancer in men. *N Engl J Med* 1999;**340**:1390-97.
- ⁸⁹ McLaughlin JK, Blot WJ, Mandel JS *et al*. Etiology of cancer of the renal pelvis. *J Natl Cancer Inst* 1983;**71**:287-91.
- ⁹⁰ Armstrong B, Doll R. Environmental factors and cancer incidence and mortality in different countries, with special reference to dietary practices. *Int J Cancer* 1975;**15**:617-31.
- ⁹¹ Morrison AS. Geographic and time trends of coffee imports and bladder cancer. *Eur J Cancer* 1978;**14**:51-54.
- ⁹² Berlin JA, Longnecker MP, Greenland S. Meta-analysis of epidemiologic dose-response data. *Epidemiology* 1993;**4**:218-28.
- ⁹³ Greenland S, Longnecker MP. Methods for trend estimation from summarized dose-response data, with applications to meta-analysis. *Am J Epidemiol* 1992;**135**:1301-09.
- ⁹⁴ Goldbohm RA, Hertog MG, Brants HA *et al*. Consumption of black tea and cancer risk: a prospective cohort study. *J Natl Cancer Inst* 1996;**88**:93-100.